


12

SOUTHERN METHODIST UNIVERSITY

AD A J 256113



DTIC
ELECTE
MAR 1 1983
S D D

DEPARTMENT OF STATISTICS

DALLAS, TEXAS 75275

DTIC FILE COPY

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

83 02 023 173

A SIMPLE APPROXIMATION FOR BIVARIATE
NORMAL PROBABILITIES

by

Robert W. Mee and D. B. Owen

Technical Report No. 169
Department of Statistics ONR Contract

December, 1982

Research sponsored by the Office of Naval Research
Contract N00014-76-C-0613

Reproduction in whole or in part is permitted
for any purpose of the United States Government

The document has been approved for
public release and sale; its distribution is unlimited

DEPARTMENT OF STATISTICS
Southern Methodist University
Dallas, Texas 75275



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

A SIMPLE APPROXIMATION FOR BIVARIATE NORMAL PROBABILITIES

by

Robert W. Mee

and

D. B. Owen

Department of Statistics
Southern Methodist University
Dallas, Texas 75275


Key Words: Bivariate Normal, Screening, Selection, Skewness, Kurtosis.

ABSTRACT

↙ The bivariate normal distribution function may be expressed as the product of a marginal normal distribution times a conditional distribution. By approximating this conditional distribution, we obtain a simple method for approximating bivariate normal probabilities. When the correlation falls in the interval $[-.5, .5]$, the maximum absolute error in our approximation is always less than .0008. The conditional distribution that we approximate is referred to as a 'normal conditioned on a truncated normal' distribution and is related to screening and selection problems.

↑

Figure 1: Comparison of pdf for Y_h and Z_h ($\rho = .7, .9, \text{ and } .95; h = 0$)



INTRODUCTION

The bivariate normal distribution is frequently employed as a model for screening procedures, where a product is either accepted or rejected based on some secondary measurement that is correlated with the performance characteristic (see, e.g., Owen, McIntire and Seymour, 1975). When screening is utilized, the proportion of all items which are rejected even though they are good and the proportion of accepted parts which are in fact defective are of interest. The purpose of this paper is to provide simple formulae for approximating these probabilities. The results that follow should also be of interest in other settings where bivariate normal probabilities are required but the potential user does not have ready access to a computer.

Tables of the bivariate normal distribution are like those of the univariate normal in that the variates are standardized, i.e., each variate has its mean subtracted and the difference is divided by the standard deviation. Hence, we start with X and Y having a joint (standardized) bivariate normal distribution with correlation ρ . We propose a simple approximation for the cumulative distribution function (cdf) $F(\cdot, \cdot; \rho)$ of (X, Y) . For any constants h and k , the probability $F(h, k; \rho)$ may be factored into a marginal probability times a conditional probability, i.e.,

$$F(h, k; \rho) = \Pr[X \leq h] \cdot \Pr[Y \leq k | X \leq h]. \quad (1)$$

We propose approximating the conditional distribution of Y , given $X \leq h$, by a normal distribution. In this way, $F(h, k; \rho)$ can be approximated as the product of two univariate normal probabilities.

Mallows (1959) obtained an approximation for $F(h, k; \rho)$ which requires simple computations and evaluation of univariate normal probabilities and

percentiles. For the values of h , k and ρ that Mallows examined, the maximum error was .0076. The approximation we present is simpler than Mallow's formulae and is more accurate unless $|\rho|$ exceeds .8. El Lozy (1982) provides a simple way to compute probabilities of the special form $F(h, \rho h; \rho)$. El Lozy's method is most accurate for ρ close to 1. In contrast, our approximation has its greatest accuracy when $|\rho|$ is less than .5.

Before we present the approximation, we investigate the conditional distribution of Y , given $X \leq h$. This distribution, which we label the normal conditioned on a truncated normal, is of interest in itself since it can represent the distribution of the performance variable in the accepted population after screening with respect to an upper specification limit.

NORMAL CONDITIONED ON A TRUNCATED NORMAL

It is well-known that the distribution of Y conditioned on $X = h$ is normal with mean ρh and variance $1 - \rho^2$. That is, Y is said to have a conditional normal distribution. However, the distribution of Y conditional on $X \leq h$ is not normal (unless $\rho = 0$), but has a probability density function (pdf) given by

$$G'(y)G[(h-\rho y)/(1-\rho^2)^{1/2}]/G(h), \quad (2)$$

where $G'(\cdot)$ and $G(\cdot)$ denote the standard normal pdf and cdf, respectively.

Let Y_h denote the random variable with pdf (2). The mean of Y_h is

$$\mu = -\rho G'(h)/G(h) \quad (3)$$

and the variance is

$$\sigma^2 = 1 + \rho h \mu - \mu^2. \quad (4)$$

Higher moments are given by

$$\mu'_n = (n-1)\mu'_{n-2} + \mu \left[\sum_{i=0}^{n-1} (1-\rho^2)^{i/2} (\rho h)^{n-1-i} \binom{h-1}{i} E(Z^i) \right] \quad (5)$$

where $\mu_n' = E(Y_h^n)$ and where $E(Z^i) = 0$ if i is odd, is equal to one if i is zero or two, and is equal to $(i-1)!2^{-(i-2)/2}[\{(i-2)/2!\}]^{-1}$ otherwise. Expressions (3) - (5) may be obtained from Johnson and Kotz (1972, pp. 112, 114).

Although the distribution of Y_h differs from the normal, unless $|\rho|$ is near one, the distribution of Y_h can be adequately approximated by a normal pdf. Let Z_h denote a normal random variable with mean given by (3) and variance given by (4). Figure 1 gives a comparison of the pdf's of Y_h and Z_h for $h = 0$ and $\rho = .7, .9$, and $.95$. We have also computed the coefficient of skewness $\gamma_1 = \mu_3/\mu_2^{3/2}$ and the measure of kurtosis $\gamma_2 = \mu_4/\mu_2^2 - 3$, where $\mu_n = E[(Y_h - \mu)^n]$. For $|\rho| \leq .5$, the pdf's of Y_h and Z_h are virtually indistinguishable. However, as $|\rho|$ approaches 1, the pdf of Y_h approaches the pdf of a normal truncated above h if $\rho > 0$ and below $-h$ if $\rho < 0$.

[Figure 1 here]

The similarity of the distribution of Y_h to a normal distribution has implications for screening problems. For example, consider the situation where an aircraft part is inspected for fractures using X-rays. Suppose lifetime (L) and maximum crack length (MCL) are modeled using a bivariate normal distribution with $\rho = -.7$. The average and standard deviation (SD) for L are taken to be 8000 hours and 1000 hours, respectively. Suppose that we reject any part with a crack exceeding 1 SD above average for MCL (i.e., we reject about 16% of the parts). Then using (3) and (4) (with $h = 1$, $\rho = -.7$), $\mu = .201$ and $\sigma = .905$ and the distribution of lifetime for accepted parts has a mean of $8000 + 1000\mu = 8201$ hours

and an SD of $1000\sigma = 905$ hours. The fifth percentile for L before selection was 6355 hours. Using the normal to approximate the distribution of L for the accepted parts, we find that, after screening, only 2% of the parts should fail before 6355 hrs. [Expressions (3) - (5) are for computing the moments of the standardized variate Y , conditional on $X \leq h$. The calculations above illustrate how to convert from μ and σ to the mean and SD in original units.]

APPROXIMATING $F(h, k; \rho)$

Visual comparison of the distributions of Y_h and Z_h suggests that, whenever $|\rho|$ is small, we may accurately approximate $F(h, k; \rho)$ by the product

$$B(h, k; \rho) = G(h) \cdot G[(k-\mu)/\sigma] \quad (6)$$

[where μ and σ are as defined in (3) and (4)], since $\Pr[Y \leq k | X \leq h] = \Pr[Y_h \leq k]$ may be approximated by $\Pr[Z_h \leq k] = G[(k-\mu)/\sigma]$.

As we investigated the accuracy of $B(h, k; \rho)$ for approximating $F(h, k; \rho)$ for various h, k and ρ , we found that the error was less when h was negative and when $|k| < |h|$. Therefore, to approximate the bivariate normal probability $F(c, d; r)$, the error of approximation is minimized by the following scheme:

1. Choose c and d so that $|c| \geq |d|$. This can always be done, since $F(c, d; r) = F(d, c; r)$.
- 2a. If $c \leq 0$, set $h = c$, $k = d$ and $\rho = r$ and approximate $F(c, d; r)$ by $B(h, k; \rho)$; or,
- 2b. If $c > 0$, set $h = -c$, $k = d$ and $\rho = -r$ and approximate $F(c, d; r) = G(d) - F(-c, d; -r)$ by $G(k) - B(h, k; \rho)$.

For example, to approximate the probability $F(.2, 1; .7)$, set $c = 1$, and following 2b, $h = -1$, $k = .2$ and $\rho = -.7$. Then $\mu = 1.0676$, $\sigma = .77946$,

$B(-1, 2; -.7) = G(-1) \cdot G[(.2-\mu)/\sigma] = .02108$, and the approximation is $G(.2) - .02108 = .55818$. The exact value is .55842, so our approximation error is only .00024.

Using this scheme, the maximum error incurred in approximating $F(c, d; r)$ for any c and d is:

$ r $	Maximum error
.1	.00001
.2	.00006
.3	.00019
.4	.00043
.5	.00079
.6	.00121
.7	.00164
.8	.00282
.9	.00765

EXAMPLE

In closing, we illustrate the usefulness of this approximation with the example of Lipow and Eidemiller (1964). Rocket motor cases are known to have a strength S that is normally distributed with mean 700 pounds/inch² (psi) and an SD of 100 psi. It is also known that peak rocket operating pressure P is normally distributed with mean 500 psi and SD 100 psi. (P and S are independently distributed.) When a rocket is fired, if the peak pressure exceeds case strength, i.e., if $\Delta = S - P$ is negative, the case will rupture and the rocket will fail. Thus, the proportion of cases that will rupture when fired is $\Pr[\Delta < 0] = G(-\sqrt{2}) = .0786$ (since

Δ is normally distributed with mean 200 psi and SD 141.42 psi). In order to decrease the frequency of failure due to rupture, the motor cases are pre-tested to a pressure of 600 psi, and discarded if they rupture. Hence, the probability of rupture for a case which passes the test is $\Pr[\Delta < 0, S \geq 600]/\Pr[S \geq 600]$. Now $\Pr[\Delta < 0, S \geq 600] = \Pr[Y < -\sqrt{2}, X \geq -1]$ with $\rho = 1/\sqrt{2}$, which also equals $F(-\sqrt{2}, 1; -\rho)$. To approximate $F(-\sqrt{2}, 1; -\rho)$, we compute

$$B(-\sqrt{2}, 1; -.7071) = G(-\sqrt{2})G[(1 - 1.319)/.761] = .0265.$$

From this bivariate probability, we can easily construct the table of probabilities:

	<u>S < 600</u>	<u>S \geq 600</u>	Totals
$\Delta < 0$.0521	.0265	.0786
$\Delta \geq 0$.1066	.8148	.9214
Totals	.1587	.8413	1.0000

Hence, the proportion of failures for the cases that passed inspection is $.0265/.8413 = .0315$. By screening, we have reduced the failure rate from 7.86% to 3.15%. However, 10.66% of the motor cases would have fired successfully, but were discarded after pre-testing.

The true value for $F(-\sqrt{2}, 1; -.7071)$ is .0267, so the error of the above approximation was .0002. This example illustrates both the simplicity and accuracy of the approximation for bivariate normal probabilities presented in this paper.

ACKNOWLEDGEMENT

This research was supported in part by the Office of Naval Research Contract No. N00014-76-C-0613.

REFERENCES

- El Lozy, M. (1982), "Simple Computation of a Bivariate Normal Integral Arising from a Problem of Misclassification with Applications to the Diagnosis of Hypertension," Communications in Statistics - Theory and Methods, Vol. A11, Issue 19, pp. 2195-2205.
- Johnson, N. L. and Kotz, S. (1972), Distributions in Statistics, 4: Multivariate Distributions. New York: John Wiley and Sons, Inc.
- Lipow, M. and Eidemilier, R. L. (1964), "Application of the Bivariate Normal Distribution to a Stress vs. Strength Problem in Reliability Analysis," Technometrics, Vol. 6, No. 3, pp. 325-328.
- Mallows, C. L. (1959), "An Approximate Formula for Bivariate Normal Probabilities," Princeton University, Statistical Techniques Research Group, Technical Report No. 30.
- Owen, D. B., McIntire, D. and Seymour E. (1975), "Tables Using One or Two Screening Variables to Increase Acceptable Product Under One-sided Specifications," Journal of Quality Technology, Vol. 7, No. 3, pp. 127-138.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 169	2. GOVT ACCESSION NO AD-A125033	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A SIMPLE APPROXIMATION FOR BIVARIATE NORMAL PROBABILITIES		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL REPORT
		6. PERFORMING ORG. REPORT NUMBER 169
7. AUTHOR(s) Robert W. Mee and D. B. Owen		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0613
9. PERFORMING ORGANIZATION NAME AND ADDRESS Southern Methodist University Dallas, Texas 75275		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 042-389
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		12. REPORT DATE December, 1982
		13. NUMBER OF PAGES 11
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited. Reproduction in whole or in part is permitted for any purposes of the United States Government.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bivariate Normal, Screening, Selection, Skewness, Kurtosis		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The bivariate normal distribution function may be expressed as the product of a marginal normal distribution times a conditional distribution. By approximating this conditional distribution, we obtain a simple method for approximating bivariate normal probabilities. When the correlation falls in the interval $[-.5, .5]$, the maximum absolute error in our ap- proximation is always less than .0008. The conditional distribution that we approximate is referred to as a 'normal conditioned on a truncated normal' distribution and is related to screening and selection problems.		

DD FORM 1473

1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601